

An Upper Bound for the Number of Planar Lattice Triangulations

Emile E. Anclin
Institute of Mathematics, MA 6-2
TU Berlin, D-10623 Berlin, Germany
anclin@math.tu-berlin.de

February 1, 2008

Abstract

We prove an exponential upper bound for the number $f(m, n)$ of all maximal triangulations of the $m \times n$ grid:

$$f(m, n) < 2^{3mn}.$$

In particular, this improves a result of S. Yu. Orevkov [1].

We consider lattice polygons P (with vertices in $\mathbb{Z}^2$), for example the convex hull of the grid $P_{m,n} := \{0, 1, \dots, m\} \times \{0, 1, \dots, n\}$. We want to estimate the number of **maximal** lattice triangulations of P , i.e., triangulations using all integer points $P \cap \mathbb{Z}^2$ in P . These are exactly the **unimodular** triangulations, in which all the triangles have integer vertices and area $\frac{1}{2}$. From now on we will talk only about unimodular triangulations. Denote by $f(P)$ the number of (unimodular) triangulations of P and by $f(n, m)$ the number of triangulations of $P_{m,n}$. S. Yu. Orevkov's upper bound [1] is $f(m, n) \leq 4^{3mn}$.

Theorem 1 *The number $f(P)$ of maximal triangulations of a lattice polygon P is bounded by*

$$f(P) \leq 2^{|E'|},$$

where $|E'|$ is the number of inner (non-boundary) edges in any unimodular triangulation of P .

In particular, the number of unimodular triangulations of the grid $P_{m,n}$ is bounded by

$$f(m, n) \leq 2^{3mn-m-n} < 2^{3mn}.$$

The Haystack Approach

Let P be a closed, not necessarily convex lattice polygon and $\text{int}(P)$ its interior. Define $M := (\frac{1}{2}\mathbb{Z}^2 \setminus \mathbb{Z}^2) \cap \text{int}(P)$, the possible midpoints of the inner edges of a lattice triangulation of P .

Lemma 2 *In any unimodular triangulation of P there is a canonical bijection between the inner edges E' and their midpoints in M .*

PROOF: The injection from E' to M is clear.

On the other hand all unimodular triangles are $SL(2, \mathbb{Z})$ -equivalent to $\mathbb{Z}^2$ -translates of $\text{conv}\{\mathbf{0}, e_1, e_2\}$, so they don't contain interior points from M . $\square$

NOTATION: For a subcomplex S of a triangulation of P and $r \in M$, if there is an edge through r in S we denote it by $e_S(r)$. We use a lexicographic order on $(\frac{1}{2}\mathbb{Z})^2$:

$$(x_1, y_1) \prec (x_2, y_2) \iff [y_1 < y_2] \text{ or } [y_1 = y_2 \text{ and } x_1 < x_2].$$

Definition 3 A **haystack** H (with respect to some $r \in M$) is a subcomplex of a triangulation of P , with an edge through $r' \in M$ if and only if $r' \prec r$.

PROOF OF THEOREM 1:

The idea is to run through M lexicographically, and at each step to add an edge through $r \in M$. We will see that in each step there are at most two possibilities to put the new edge through r .




Figure 1: A haystack with respect to r .

We proceed by induction on the totally ordered set $(M, \prec)$, thus proving that the number of haystacks with respect to some $r \in M$ is $\leq 2^{e_r}$ where e_r is the number of predecessors of r in M . Thus after the final step (that is, after processing the largest r in $(M, \prec)$) we have obtained that there are at most $2^{|M|} = 2^{|E'|}$ unimodular triangulations of P .

Now for some $r \in M$ consider a haystack H with respect to r . We want to add a “needle” to our haystack so that the resulting subcomplex will again be a haystack. So we consider the set A_r of possible endpoints v of edges through r , with $v \prec r$:

$$A_r := \{v \in \mathbb{Z}^2 \mid v \prec r \text{ and } H \cup \{[v, v + 2\vec{r}]\} \text{ is a haystack}\}.$$

We want to prove that $|A_r| \leq 2$ for all $r \in M$.

We say that v is **visible** from r if the edge $[v, r]$ crosses no other edge or integral point. Consider

$$A := \{v \in \text{conv}(\{r\} \cup A_r) \cap \mathbb{Z}^2 \mid v \text{ is visible from } r\}.$$

As $v \in A_r$ is visible from r we have $A \supseteq A_r$. Furthermore $v \prec r$ holds for all $v \in A$. (See Figure 2).




Figure 2: Here $A_r = \{v_1, v_3\}$ and $A = \{v_1, v_2, v_3\}$, while by definition $v, v' \notin A$ and $\alpha_1 < \alpha_2 < \alpha_3$.

We now order A by the angles $\alpha(v)$ of $r\vec{v}$ with the x -axis turning counter-clockwise and starting by π , so that we have $\alpha_i = \alpha(v_i)$, $\alpha_1 < \alpha_2 < \dots < \alpha_k$. Indeed, we never have $\alpha_i = \alpha_j$, otherwise r, v_i, v_j would lie on a line, but then one of the two points v_i, v_j would not be visible from r , because both are $\prec r$.


Observe that $v_1 \in A_r$: $v \prec r$ for all $v \in A$, so a point v with a smaller angle to the x -axis than the first one in A_r can't be in $\text{conv}(A_r \cup \{r\}) \supset A$.

We say that a triangle $[v_i, v_{i+1}, r]$ is **empty** if there is no edge through it and no $\frac{1}{2}\mathbb{Z}^2$ -points in its interior. The triangle $[v_i, v_{i+1}, r]$ is empty as A contains all points in $\text{conv}(A_r \cup \{r\}) \supset [v_i, v_{i+1}, r]$ visible from r . The midpoint $s_i := \frac{1}{2}(v_i + v_{i+1})$ is half-integer, $s_i \in M$, as $[v_i, v_{i+1}, r]$ is empty. We also have $s_i \prec r$, and so $e_H(s_i) = [v_i, v_{i+1}]$, since the triangle $[v_i, v_{i+1}, r]$ is empty. Additionally it has area $\frac{1}{4}$, otherwise the triangle wouldn't be empty.

Define $w_i := r + v_i\vec{r}$ and $r' := \frac{1}{2}(v_1 + w_2)$, $r'' := \frac{1}{2}(v_2 + w_1)$. Then v_1, w_2, v_2, w_1 form a parallelogram with center r , and r, r', r'' are on a line (parallel to (v_1v_2)). So either $r' \prec r$ or $r'' \prec r$.

Case 1: Suppose first that $r' \prec r$.

The triangle $\Delta = [v_1, v_2, w_2]$ is unimodular as $\text{area}(\Delta) = 2 \text{area}[v_1, v_2, r] = \frac{1}{2}$; so there are no integer points between the line (w_1w_2) and the line (v_1v_2) . The edge $e_H(r')$ has nonempty intersection with these two lines (but doesn't cross $[v_1, w_1]$, since $v_1 \in A_r$).



But where could a third point $v \in A_r$ (other than v_1, v_3) be? The line $(r'r)$ is parallel to (v_1v_2) , we have $\alpha(r') < \alpha_i \leq \alpha_i$; and $r' \prec r$, $v \prec r$ for all $v \in A_r$. So all points of A are on the same side of $(r'r)$ as v_1 and v_2 . So v is on or beyond the line (v_1v_2) and hence the

edge through r starting at v would necessarily cross the edge $e_H(r')$. So there can be no other point v in A_r , that is, $|A_r| \leq 2$.

Case 2: The situation for $r'' \prec r$ is similar:

The edge through r'' must be $e(r'') = [v_2, w_1]$, otherwise it would cut $[v_1, w_1]$ or $[v_1, v_2]$; in the first case we would have $v_1 \notin A_r$ and in the second case v_2 wouldn't be visible from r . And $[v_1, v_2, w_1]$ is again unimodular, so there is no possibility for a third $v \in A_r$. $\square$

Our Theorem 1 and its proof clearly extend to a more general situation, namely the case of a not necessarily simply connected lattice polygon (which may have holes), possibly with additional, fixed inner edges.

We can define the capacities $c_{m,n} := \frac{\log_2 f(m,n)}{mn}$; see [2]. From sublinearity of $f(m,n)$ it follows by Fekete's lemma [3, p. 85] that the limit capacities

$$c_m := \lim_{n \rightarrow \infty} \frac{\log_2 f(m,n)}{mn}, \quad c_\Delta := \lim_{n \rightarrow \infty} \frac{\log_2 f(n,n)}{n^2}$$

exist. Theorem 1 yields the upper bounds

$$c_m \leq 3 - \frac{1}{m},$$

which includes the best known upper bounds for all c_m (compare [2]).

In generating triangulations with the “haystack approach” as in the proof of Theorem 1, one will in many situations have $|A_r| = 1$. So probably our upper bound $c_\Delta \leq 3$ for the limit capacity c_Δ is not sharp.

As for lower bounds, the recursion formulas for $(n \times 2)$ - and $(n \times 3)$ -strips, as given in [2], together with submultiplicativity, show that $c_\Delta > 2.055$.

References

- [1] S. YU. OREVKOV: Asymptotic number of triangulations with vertices in $\mathbb{Z}^2$, *J. Combinatorial Theory, Ser. A* **86** (1999), 200-203.
- [2] V. KAIBEL & G. M. ZIEGLER: Counting unimodular lattice triangulations, Preprint, TU Berlin, November 2002, 31 pages; [arXiv:math.CO/0211268](#); to appear in *British Combinatorial Surveys* (C. D. Wensley, ed.), Cambridge University Press 2003.
- [3] J. H. VAN LINT & R. M. WILSON: A Course in Combinatorics, *Cambridge University Press*, Cambridge 1992.